

Fig. 1.

		mortality	%	anaesth	technical
	<b>Controls:</b>	<b>11/39</b>	<b>(28,2)</b>	<b>3</b>	<b>8</b>
	<b>IL-6:</b>	<b>10/80</b>	<b>(12,5)</b>	<b>5</b>	<b>5</b>
<b>L.D.</b> (Low Doses)	IL 0,1:	0/12	-	-	-
	IL 1:	2/14	(14,2)	2	-
	IL 10:	2/20	(10,0)	2	-
<b>H.D.</b> (High Doses)	IL 100:	4/21	(19,0)	1	3
	IL 500:	2/13	(15,4)	0	2
	<b>IL L.D.:</b>	<b>4/46</b>	<b>( 8,7)</b>		
	<b>IL H.D.:</b>	<b>6/34</b>	<b>(17,6)</b>		

Fig. 2




Fig. 3





Fig. 4

Hours after injection (100µg/Kg):

STAT-3


AP-1

NFkB



BEST AVAILABLE COPY


Fig. 5



BEST AVAILABLE COPY

Fig 6

IL-6 receptors in normal and cirrhotic rats: gp 130 and gp 80



BEST AVAILABLE COPY

Fig. 7




Fig. 8







Fig. 9



BEST AVAILABLE COPY